

A Maillet type theorem for generalized power series

R. R. Gontsov

Let us consider an ordinary differential equation (ODE)

$$F(z, u, \delta u, \dots, \delta^m u) = 0 \quad (1)$$

of order m with respect to the unknown u , where $F(z, u_0, u_1, \dots, u_m) \neq 0$ is a polynomial of $m + 2$ variables, $\delta = z \frac{d}{dz}$.

The classical Maillet theorem [1] asserts that any formal power series solution $\varphi = \sum_{n=0}^{\infty} c_n z^n \in \mathbb{C}[[z]]$ of (1) is a power series of Gevrey order $1/k$ for some $k \in \mathbb{R}_{>0} \cup \{\infty\}$. This means that the power series

$$f = \sum_{n=0}^{\infty} \frac{c_n}{\Gamma(1 + n/k)} z^n$$

converges in a neighbourhood of zero, where Γ is the Euler gamma-function. In other words,

$$|c_n| \leqslant A B^n (n!)^{1/k}$$

for some $A, B > 0$.

First exact estimates for the Gevrey order of a power series formally satisfying an ODE were obtained by J.-P. Ramis [2] in the linear case,

$$Lu = a_m(z)\delta^m u + a_{m-1}(z)\delta^{m-1}u + \dots + a_0(z)u = 0, \quad a_i \in \mathbb{C}\{z\}.$$

He has proved that such a power series is of the *exact* Gevrey order $1/k \in \{0, 1/k_1, \dots, 1/k_s\}$, where $k_1 < \dots < k_s < \infty$ are all of the positive slopes of the Newton polygon $\mathcal{N}(L)$ of the operator L . This polygon is defined as the boundary curve of a convex hull of a union $\bigcup_{i=0}^m X_i$,

$$X_i = \{(x, y) \in \mathbb{R}^2 \mid x \leqslant i, y \geqslant \text{ord}_0 a_i(z)\}, \quad i = 0, 1, \dots, m$$

(see Fig. 1). The exactness of the Gevrey order means that there is no $k' > k$ such that the power series is of the Gevrey order $1/k'$.




Figure 1: The Newton polygon $\mathcal{N}(L)$ with two positive slopes $k_1 = (\text{ord}_0 a_j - \text{ord}_0 a_i)/(j - i)$, $k_2 = (\text{ord}_0 a_m - \text{ord}_0 a_j)/(m - j)$.

The result of Ramis has been further generalized by B. Malgrange and Y. Sibuya for a non-linear ODE of the general form (1).

Theorem 1 (Malgrange [3]). *Let $\varphi \in \mathbb{C}[[z]]$ satisfy the equation (1), that is $F(z, \Phi) = 0$, where $\Phi = (\varphi, \delta\varphi, \dots, \delta^m\varphi)$, and $\frac{\partial F}{\partial u_m}(z, \Phi) \neq 0$. Then φ is a power series of Gevrey order $1/k$, where k is the least of all the positive slopes of the Newton polygon $\mathcal{N}(L_\varphi)$ of a linear operator*

$$L_\varphi = \sum_{i=0}^m \frac{\partial F}{\partial u_i}(z, \Phi) \delta^i$$

(or $k = +\infty$, if $\mathcal{N}(L_\varphi)$ has no positive slopes).

The refinement of Theorem 1 belongs to Sibuya [4, App. 2]. This claims that φ is a power series of the *exact* Gevrey order $1/k \in \{0, 1/k_1, \dots, 1/k_s\}$, where $k_1 < \dots < k_s < \infty$ are all of the positive slopes of the Newton polygon $\mathcal{N}(L_\varphi)$.

In the talk we study *generalized* power series solutions of (1) of the form

$$\varphi = \sum_{n=0}^{\infty} c_n z^{s_n}, \quad c_n \in \mathbb{C}, \quad s_n \in \mathbb{C}, \quad (2)$$

with the power exponents satisfying conditions

$$0 \leq \operatorname{Re} s_0 \leq \operatorname{Re} s_1 \leq \dots, \quad \lim_{n \rightarrow \infty} \operatorname{Re} s_n = +\infty$$

(the latter, in particular, implies that a set of exponents having a fixed real part is finite).

For the generalized power series (2) one may naturally define the *valuation*

$$\operatorname{val} \varphi = s_0,$$

and this is also well defined for any polynomial in $z, \varphi, \delta\varphi, \dots, \delta^m \varphi$.

The main result of the present talk is an analog of the Maillet theorem (more precisely, of the Malgrange theorem) for generalized power series.

Theorem 2 ([5]). *Let the generalized power series (2) formally satisfy the equation (1), $\frac{\partial F}{\partial u_m}(z, \Phi) \neq 0$, and for each $i = 0, 1, \dots, m$ one have*

$$\frac{\partial F}{\partial u_i}(z, \Phi) = A_i z^\lambda + B_i z^{\lambda_i} + \dots, \quad \operatorname{Re} \lambda_i > \operatorname{Re} \lambda, \quad (3)$$

where not all A_i equal zero. Let k be the least of all the positive slopes of the Newton polygon $\mathcal{N}(L_\varphi)$ (or $k = +\infty$, if $\mathcal{N}(L_\varphi)$ has no positive slopes). Then for any sector S of sufficiently small radius with the vertex at the origin and of the opening less than 2π , the series

$$\sum_{n=0}^{\infty} \frac{c_n}{\Gamma(1 + s_n/k)} z^{s_n}$$

converges uniformly in S .

Remark 1. The Newton polygon of L_φ in the case of the generalized power series φ is defined similarly to the classical case: this is the boundary curve of a convex hull of a union

$$\bigcup_{i=0}^m \left\{ (x, y) \in \mathbb{R}^2 \mid x \leq i, y \geq \operatorname{Re} \operatorname{val} \frac{\partial F}{\partial u_i}(z, \Phi) \right\}.$$

Note that $k = +\infty$ in Theorem 2 if and only if $A_m \neq 0$. In this case φ does converge in S , which has been already proved in [6] by the majorant method.

Remark 2. From Theorem 2 one deduces the following estimate for the coefficients c_n of the formal series solution (2) of (1):

$$|c_n| \leq A B^{\operatorname{Re} s_n} |\Gamma(1 + s_n/k)|,$$

for some $A, B > 0$.

Example 1. We consider an equation

$$z^3 u u'' - z u' + \sqrt{2} u + z^2 = 0,$$

or, written with use of the operator δ ,

$$z u \delta^2 u - (z u + 1) \delta u + \sqrt{2} u + z^2 = 0.$$

It has a generalized power series solution of the form $\varphi = \sum_{s \in K} c_s z^s$, where

$$K = \{\sqrt{2} + m_1(1 + \sqrt{2}) + m_2(2 - \sqrt{2}) \mid m_1, m_2 \in \mathbb{Z}_+\}.$$

The first terms of φ are

$$\varphi = z^{\sqrt{2}} + \left(1 + \frac{\sqrt{2}}{2}\right)z^2 + (3\sqrt{2} - 4)z^{2\sqrt{2}+1} + \dots$$

Since $F(z, u_0, u_1, u_2) = zu_0u_2 - (zu_0 + 1)u_1 + \sqrt{2}u_0 + z^2$, we have

$$\begin{aligned} \frac{\partial F}{\partial u_0}(z, \Phi) &= z\delta^2\varphi - z\delta\varphi + \sqrt{2} = \sqrt{2} + \dots, \\ \frac{\partial F}{\partial u_1}(z, \Phi) &= -z\varphi - 1 = -1 + \dots, \\ \frac{\partial F}{\partial u_2}(z, \Phi) &= z\varphi = z^{\sqrt{2}+1} + \dots \end{aligned}$$

Therefore, the Newton polygon $\mathcal{N}(L_\varphi)$ is determined by the points $(0, 0)$, $(1, 0)$, and $(2, 1 + \sqrt{2})$. Hence, $k = 1 + \sqrt{2}$ is a unique positive slope of $\mathcal{N}(L_\varphi)$ and, by Theorem 2,

$$|c_s| \leq AB^s \Gamma\left(1 + \frac{s}{1 + \sqrt{2}}\right),$$

for some $A, B > 0$.

References

- [1] E. Maillet, Sur les séries divergentes et les équations différentielles, *Ann. Sci. Ecole Norm. Sup.* **3** (1903), 487–518.
- [2] J.-P. Ramis, Dévissage Gevrey, *Astérisque* **59/60** (1978), 173–204.
- [3] B. Malgrange, Sur le théorème de Maillet, *Asympt. Anal.* **2** (1989), 1–4.
- [4] Y. Sibuya, Linear Differential Equations in the Complex Domain: Problems of Analytic Continuation, *Transl. Math. Monographs* **82**, A.M.S., 1990.
- [5] R. R. Gontsov, I. V. Goryuchkina, The Maillet–Malgrange type theorem for generalized power series, arXiv:1601.05778, 2016, 9pp.
- [6] R. R. Gontsov, I. V. Goryuchkina, On the convergence of generalized power series satisfying an algebraic ODE, *Asympt. Anal.* **93**:4 (2015), 311–325.